

Interleukin-10 Protein

Cat.No:DTD0062

DESCRIPTION

Mana	laterie do anteir
Name	Interleukin-10 protein
Purification	The soluble target protein Interleukin-10 was derived from secreted expression and
	obtained by affinity chromatography purification
Protein	At 0.400 mg/ml, BSA was used as the standard and the protein concentration was
Concentration	determined by BCA
Molecular weight	48 kDa
Purity	>90% by SDS-PAGE
Species	Dog
Expression	HEK293
System	
Endotoxin	<1EU / mg, as determined by a gel-based endotoxin detection kit
Physical	Liquid
Appearance	
Buffer	1×PBS, pH7.4
Stability & Storage	For long term use, aliquot and store at -80°C (avoiding multiple freeze-thaws)
SDS-PAGE & Western Blot	kDa 1 M1 2 kDa M2 2
	180 120
	120 80 Lane 1: BSA (1.00µg)
	80 80 Lane 1: BSA (1.00μg) 65 65 Lane 2: Interleukin-10 protein(0.80μg)
	50 50 M ₁ : SDS-PAGE Marker
	40 40 Mo: Western Blot Marker
	30 30 Using Anti-His antibody
	20 15 10 20
	SDS-PAGE Western Blot

Thank you for choosing Detai Biological. We are dedicated to serving you! For Research Use Only