

Interleukin-10 Protein

Cat.No:DTD0082

DESCRIPTION

Name	Interleukin-10 protein
Purification	The soluble target protein Interleukin-10 was derived from secreted expression and obtained by affinity chromatography purification
Protein Concentration	At 0.890 mg/ml, BSA was used as the standard and the protein concentration was determined by BCA
Molecular weight	113 kDa
Purity	>85% by SDS-PAGE
Species	Human
Expression System	ExpiCHO
Physical Appearance	Liquid
Buffer	1×PBS,5 % Glycerol,pH7.4
Stability & Storage	For long term use, aliquot and store at -80°C (avoiding multiple freeze-thaws)
SDS-PAGE & Western Blot	M1 1 2 M2 2 kDa kDa Lane 1: BSA (1.00 µg) Lane 2: Interleukin-10-HSA-IgG Fc 120 120 80 Lane 2: Interleukin-10-HSA-IgG Fc 50 65 50 M1: SDS-PAGE Marker 40 30 30 M2 M2 20 50 65 M2 M2 50 30 20 M2 M2 50 30 20 M2 M2 50 50 40 30 M2 20 30 20 M2 M2 50 20 30 20 M2 SDS-PAGE Western Blot M2 M2
	SDS-PAGE Western Blot

Thank you for choosing Detai Biological. We are dedicated to serving you! For Research Use Only