

## Ras-related C3 botulinum toxin substrate 2 Protein

Cat.No:DTD0129

## **DESCRIPTION**

Name	Ras-related C3 botulinum toxin substrate 2 protein
Purification	The soluble target protein Ras-related C3 botulinum toxin substrate 2 was derived from secreted expression and obtained by affinity chromatography purification
Protein	At 0.400 mg/ml, BSA was used as the standard and the protein concentration was
Concentration	determined by BCA
Molecular weight	37 kDa
Purity	>85% by SDS-PAGE
Expression System	HEK293
Species	Human
Physical Appearance	Liquid
Buffer	1×PBS, pH7.4
Stability & Storage	For long term use, aliquot and store at -80°C (avoiding multiple freeze-thaws)
SDS-PAGE & Western Blot	kDa M₁ 1 2 kDa M₂ 2  180 120 120 120 120 80 65 50 40 30 20 15 10  kDa M₂ 2  Lane 1: BSA (1.00 μg) Lane 2: Ras-related C3 botulinum toxin substrate 2 protein (0.80μg) M₁: SDS-PAGE Marker M₂: Western Blot Marker Using Anti-His antibody
	SDS-PAGE Western Blot

Thank you for choosing Detai Biological. We are dedicated to serving you!

For Research Use Only

Tel: 400-626-9766 www.DetaiBio.com Sales@DetaiBio.com